

Lawrence Berkeley National Laboratory

Recent Work

Title

Longitudinal fluctuations and decorrelations of anisotropic flows in relativistic heavy-ion collisions

Permalink

<https://escholarship.org/uc/item/0sh195wm>

Authors

Wu, XY
Pang, LG
Qin, GY
et al.

Publication Date

2019-02-01

DOI

10.1016/j.nuclphysa.2018.09.014

Peer reviewed



Longitudinal fluctuations and decorrelations of anisotropic flows in relativistic heavy-ion collisions

Xiang-Yu Wu^a, Long-Gang Pang^{b,c}, Guang-You Qin^a, Xin-Nian Wang^{a,b,c}

^a*Institute of Particle Physics and Key Laboratory of Quark and Lepton Physics (MOE), Central China Normal University, Wuhan, Hubei 430079, China*

^b*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA*

^c*Physics Department, University of California, Berkeley, CA 94720, USA*

Abstract

We study the longitudinal decorrelations of elliptic, triangular and quadrangular flows in heavy-ion collisions at the LHC and RHIC energies. The event-by-event CLVisc (3+1)-dimensional hydrodynamics model, combined with the fully fluctuating AMPT initial conditions, is utilized to simulate the space-time evolution of the strongly-coupled quark-gluon plasma. Detailed analysis is performed for the longitudinal decorrelations of flow vectors, flow magnitudes and flow orientations. We find strong correlations between final-state longitudinal decorrelations of anisotropic flows and initial-state longitudinal structures and collision geometry: while the decorrelation of elliptic flow shows a non-monotonic centrality dependence due to initial elliptic geometry, typically the longitudinal flow decorrelations are larger in lower energy and less central collisions where the mean lengths of the string structure are shorter in the initial states.

Keywords: quark-gluon plasma, anisotropic flows, fluctuations, longitudinal decorrelations, relativistic hydrodynamics

1. Introduction

Initial-state fluctuations and final-state anisotropic flows V_n provide important tools for studying strongly-coupled quark gluon plasma (QGP) in relativistic nuclear collisions at RHIC and the LHC. By means of relativistic hydrodynamics simulation supplemented with fluctuating initial conditions, many theoretical studies have been focus on anisotropic flows and their fluctuations and correlations in the transverse directions. Recently, much interest has been paid on the fluctuations in the longitudinal (pseudorapidity) direction, which have the potential probe the longitudinal structure and evolution dynamics of the QGP fireball. Longitudinal fluctuations in the initial states imply that the final anisotropic flows may be different at two different rapidities: $V_n(\eta_1) \neq V_n(\eta_2)$ for $\eta_1 \neq \eta_2$. In this work [1], we employ the CLVisc (ideal) (3+1)-dimensional hydrodynamics model [2] combined with the AMPT [3] initial conditions, and perform a systematic study on the longitudinal decorrelations of anisotropic flows, in terms of flow vectors, flow magnitudes and flow orientations, for different collision energies and centralities in heavy-ion collisions at RHIC and the LHC.

2. Event-by-event (3+1)-dimensional hydrodynamics simulation

In this work, we use the ideal version of the CLVisc (3+1)-dimensional hydrodynamics model [2] to simulate the dynamical evolution of the QGP fireball. The initial conditions are obtained by the AMPT




Fig. 1. The slope parameters $f[n,k]$, $f_\Phi[n,k]$ and $f_M[n,k]$ for the longitudinal decorrelation functions $r[n,k](\eta)$, $r_\Phi[n,k](\eta)$ and $r_M[n,k](\eta)$ with $n=2$ (left), $n=3$ (middle), $n=4$ (right) and $k=1$, as a function of collision centrality (N_{part}) for Pb+Pb collisions at 5.02A TeV (upper), 2.76A TeV (middle) and for Au+Au collisions at 200A GeV (lower).

model [3] which provides partons' positions and momenta for constructing the energy-momentum tensor:

$$T^{\mu\nu}(\tau_0, x, y, \eta_s) = K \sum_i \frac{p_i^\mu p_i^\nu}{p_i^\tau} \frac{1}{\tau_0} \frac{1}{\sqrt{2\pi\sigma_{\eta_s}^2}} \frac{1}{2\pi\sigma_r^2} \exp \left[-\frac{(x-x_i)^2 + (y-y_i)^2}{2\sigma_r^2} - \frac{(\eta_s - \eta_{si})^2}{2\sigma_{\eta_s}^2} \right] \quad (1)$$

The ideal hydrodynamic equation $[\partial_\mu T^{\mu\nu} = 0, T^{\mu\nu} = U^\mu U^\nu (e + P) - P g^{\mu\nu}]$ is then numerically solved using KT algorithm. The momentum distributions of final hadrons are computed using Copper-Frye formula, in which the freeze-out temperature is taken as $T_{fz} = 137\text{MeV}$.

3. Longitudinal decorrelations of anisotropic flows V_n

To study anisotropic flows, we use \mathbf{Q}_n -vector method: $\mathbf{Q}_n(\eta) = \frac{1}{N} \sum_{i=1}^N e^{in\phi_i} = q_n(\eta) \hat{Q}_n(\eta) = q_n(\eta) e^{in\Phi_n(\eta)}$, where q_n and Φ_n denote flow magnitude and flow orientation, respectively, in a given η bin. The longitudinal decorrelations of anisotropic flows may be studied using the following correlation functions [4, 5, 6, 7, 8]:

$$r[n,k](\eta) = \frac{\langle \mathbf{Q}_n^k(-\eta) \mathbf{Q}_n^{*k}(\eta_r) \rangle}{\langle \mathbf{Q}_n^k(\eta) \mathbf{Q}_n^{*k}(\eta_r) \rangle}, \quad r_M[n,k](\eta) = \frac{\langle q_n^k(-\eta) q_n^k(\eta_r) \rangle}{\langle q_n^k(\eta) q_n^k(\eta_r) \rangle}, \quad r_\Phi[n,k](\eta) = \frac{\langle \hat{Q}_n^k(-\eta) \hat{Q}_n^{*k}(\eta_r) \rangle}{\langle \hat{Q}_n^k(\eta) \hat{Q}_n^{*k}(\eta_r) \rangle}, \quad (2)$$

where $r[n,k](\eta)$, $r_M[n,k](\eta)$ and $r_\Phi[n,k](\eta)$ quantify the longitudinal decorrelations of flow vectors, flow magnitudes, and flow orientations, respectively. Since the longitudinal decorrelation functions $r[n,k](\eta)$, $r_M[n,k](\eta)$ and $r_\Phi[n,k](\eta)$ decrease almost linearly as a function of pseudorapidity η , especially in small η




Fig. 2. The string structures formed by the initial patrons in the AMPT model for typical events in Pb+Pb collisions at 5.02A TeV (left) and 2.76A TeV (middle) and for Au+Au collisions at 200A GeV (right).

region [9, 10], it is convenient to use the slopes of the decorrelation functions to quantify the longitudinal decorrelation effects. We follow ATLAS Collaboration [5] to define slope parameters as follows:

$$f[n, k] = \frac{\sum_i \{1 - r[n, k](\eta_i)\} \eta_i}{2 \sum_i \eta_i^2}, f_M[n, k] = \frac{\sum_i \{1 - r_M[n, k](\eta_i)\} \eta_i}{2 \sum_i \eta_i^2}, f_\Phi[n, k] = \frac{\sum_i \{1 - r_\Phi[n, k](\eta_i)\} \eta_i}{2 \sum_i \eta_i^2}. \quad (3)$$

Figure 1 shows the slope parameters $f[n, k]$, $f_\Phi[n, k]$ and $f_M[n, k]$ (with $k = 1$), as function of collision centrality, for elliptic ($n=2$), triangular ($n=3$) and quadrangular ($n=4$) flows, for 2.76 TeV and 5.02 TeV Pb+Pb and 200 GeV Au+Au collisions. One can see that for all collision energies and centralities explored here, pure flow orientations have larger decorrelation effects than pure flow magnitudes, while the decorrelations of flow vectors sit in the middle. Also we observe a strong and non-monotonic centrality dependence for V_2 decorrelation due to the initial-state elliptic collision geometry. Since V_3 is dominated by fluctuations [11], we observe weak centrality dependence for its longitudinal decorrelation, which tends to increase from central to less central collisions (similar trend for V_4 decorrelation as well). Another important result is that V_n decorrelations are larger at RHIC (lower collision energy) than at the LHC (higher collision energies).

The collision energy and centrality dependences of longitudinal decorrelations can be traced back to the longitudinal structures in the initial states. To illustrate this, we apply k-means algorithm to extract the lengths of the initial string structures in the AMPT model. The results are shown in Figure 2, for three typical central events in 200A GeV Au+Au collisions and 2.76A TeV, 5.02A TeV Pb+Pb collisions. We can see that the mean lengths of the initial string structures increase from lower to higher collision energies. Figure 3 shows the slope parameters $f[n, k]$, $f_\Phi[n, k]$ and $f_M[n, k]$ (with $k = 1$) as function of collision energy and centrality (in terms of the mean string length) for V_2 , V_3 and V_4 at RHIC and the LHC energies. We find strong correlations between the longitudinal flow decorrelations in the final states and the longitudinal structures in the initial states: the longitudinal decorrelations of V_n tend to be larger in lower energy and less central heavy-ion collisions where the lengths of the initial string structures are shorter.

4. Summary

In summary, we have performed a detailed analysis on the longitudinal decorrelations of elliptic, triangular and quadrangular flows in relativistic heavy-ion collisions at the LHC and RHIC energies. We find that the longitudinal decorrelation effects are much larger for pure flow orientations than for pure flow magnitudes. Also, a non-monotonic centrality dependence is found for the elliptic flow decorrelation. Another important finding is that the longitudinal flow decorrelations are usually larger in lower energy and less central collisions due to the shorter lengths of the string structures in the initial states of heavy-ion collisions.

Acknowledgments

This work is supported in part by NSFC (Nos. 11775095, 11375072 and 11221504), by the Major State Basic Research Development Program in China (No. 2014CB845404), by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, DNP, of the U.S. DOE (No. DE-AC02-05CH1123), and by the U.S. NSF within the framework of the JETSCAPE collaboration (No. ACI-1550228).




Fig. 3. The slope parameters $f[n, k]$ (upper), $f_\phi[n, k]$ (middle) and $f_M[n, k]$ (lower) for the longitudinal decorrelation functions $r[n, k](\eta)$, $r_\phi[n, k](\eta)$ and $r_M[n, k](\eta)$ with $n=2$ (left), $n=3$ (middle), $n=4$ (right) and $k=1$ as function of collision energy and centrality (using the mean string length $\langle I_s \rangle$) for Pb+Pb collisions at 5.02A TeV and 2.76A TeV and for Au+Au collisions at 200A GeV. In each panel, three curves denote three collision energies: 200A GeV, 2.76A TeV and 5.02A TeV, from left to right. For each curve, five symbols represent five centrality bins: 40-50%, 30-40%, 20-30%, 10-20% and 5-10%, from left to right.

References

- [1] X.-Y. Wu, L.-G. Pang, G.-Y. Qin, X.-N. Wang, Longitudinal fluctuations and decorrelations of anisotropic flows at the LHC and RHIC energies [arXiv:1805.03762](#).
- [2] L.-G. Pang, H. Petersen, X.-N. Wang, Pseudorapidity distribution and decorrelation of anisotropic flow within CLVisc hydrodynamics [arXiv:1802.04449](#).
- [3] Z.-W. Lin, C. M. Ko, B.-A. Li, B. Zhang, S. Pal, A Multi-phase transport model for relativistic heavy ion collisions, *Phys. Rev. C* **72** (2005) 064901. [arXiv:nuc1-th/0411110](#), doi:10.1103/PhysRevC.72.064901.
- [4] V. Khachatryan, et al., Evidence for transverse momentum and pseudorapidity dependent event plane fluctuations in PbPb and pPb collisions, *Phys. Rev. C* **92** (3) (2015) 034911. [arXiv:1503.01692](#), doi:10.1103/PhysRevC.92.034911.
- [5] M. Aaboud, et al., Measurement of longitudinal flow decorrelations in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ and 5.02 TeV with the ATLAS detector, *Eur. Phys. J. C* **78** (2) (2018) 142. [arXiv:1709.02301](#), doi:10.1140/epjc/s10052-018-5605-7.
- [6] J. Jia, P. Huo, A method for studying the rapidity fluctuation and decorrelation of harmonic flow in heavy-ion collisions, *Phys. Rev. C* **90** (3) (2014) 034905. [arXiv:1402.6680](#), doi:10.1103/PhysRevC.90.034905.
- [7] J. Jia, P. Huo, Forward-backward eccentricity and participant-plane angle fluctuations and their influences on longitudinal dynamics of collective flow, *Phys. Rev. C* **90** (3) (2014) 034915. [arXiv:1403.6077](#), doi:10.1103/PhysRevC.90.034915.
- [8] P. Bozek, W. Broniowski, Longitudinal decorrelation measures of flow magnitude and event-plane angles in ultrarelativistic nuclear collisions, *Phys. Rev. C* **97** (3) (2018) 034913. [arXiv:1711.03325](#), doi:10.1103/PhysRevC.97.034913.
- [9] L.-G. Pang, G.-Y. Qin, V. Roy, X.-N. Wang, G.-L. Ma, Longitudinal decorrelation of anisotropic flows in heavy-ion collisions at the CERN Large Hadron Collider, *Phys. Rev. C* **91** (4) (2015) 044904. [arXiv:1410.8690](#), doi:10.1103/PhysRevC.91.044904.
- [10] L.-G. Pang, H. Petersen, G.-Y. Qin, V. Roy, X.-N. Wang, Decorrelation of anisotropic flow along the longitudinal direction, *Eur. Phys. J. A* **52** (4) (2016) 97. [arXiv:1511.04131](#), doi:10.1140/epja/i2016-16097-x.
- [11] G.-Y. Qin, H. Petersen, S. A. Bass, B. Muller, Translation of collision geometry fluctuations into momentum anisotropies in relativistic heavy-ion collisions, *Phys. Rev. C* **82** (2010) 064903. [arXiv:1009.1847](#), doi:10.1103/PhysRevC.82.064903.